

ECT*, Villa Tambosi, I-38050 Villazzano (TN), INFN G.C. Trento, Italy Dmitri N. Voskresensky GSI,
Planck Str. 1, D-64291 Darmstadt, Germany Moscow Engineering Physical Institute, Kashirskoe shosse 31,
RU-115409 Moscow, RussiaPermanent address

abstract Kaon polarization operator in dense baryonic matter of arbitrary isotopic composition is calculated including s- and p-wave kaon-baryon interactions. The regular part of the polarization operator is extracted from the realistic kaon-nucleon interaction based on the chiral and $1/N_c$ expansion. Contributions of the $\Lambda(1116)$, $\Sigma(1195)$, $\Sigma^*(1385)$ resonances are taken explicitly into account in the pole and regular terms with inclusion of mean-field potentials. The baryon-baryon correlations are incorporated and fluctuation contributions are estimated. Results are applied for K^- in neutron star matter. Within our model a second-order phase transition to the s-wave K^- condensate state occurs at $\rho_c \sim 4\rho_0$ once the baryon-baryon correlations are included. We show that the second-order phase transition to the p-wave K^- condensate state may occur at densities $\rho_c \sim 3 \div 5\rho_0$ in dependence on the parameter choice. We demonstrate that a first-order phase transition to a proton-enriched (approximately isospin-symmetric) nucleon matter with a p-wave K^- condensate can occur at smaller densities, $\rho < \sim 2\rho_0$. The transition is accompanied by the suppression of hyperon concentrations.